

Document Detail

Type: PMF IBTS SPEC

Document No.: IBTS/PMF/SPEC/0232[4]

Title: RED CELLS, SUITABLE FOR NEONATAL USE FOR 5 DAYS

AFTER DATE DRAWN

Owner: QA DOC CON QA DOC CONTROL

Status CURRENT
Effective Date: 13-May-2021
Expiration Date: 04-Aug-2025

Review

Review: IBTS PMF REVIEW

Level	Owner Role	Actor	Sign-off By
1	DOCUMENT CONTROLLER	REBECCA WALDEN	REBECCA WALDEN
2	QUALITY ASSURANCE WRITER IBTS	REBECCA WALDEN	REBECCA WALDEN
3	LABS PHS DIR IBTS	BARRY DOYLE	BARRY DOYLE
3	NATIONAL MEDICAL DIRECTOR	STEPHEN FIELD	STEPHEN FIELD
4	QUALITY ASSURANCE REVIEWER IBTS	COLIN JOHNS	COLIN JOHNS

Review: IBTS DOC PERIODIC REVIEW

f By

1	DOCUMENT CONTROLLER	REBECCA WALDEN	REBECCA WALDEN
2	QUALITY ASSURANCE WRITER IBTS	REBECCA WALDEN	REBECCA WALDEN
3	NATIONAL MEDICAL DIRECTOR	TOR HERVIG	TOR HERVIG
3	COMPONENTS HEAD OF DEPT MRTC	AINE FITZPATRICK	AINE FITZPATRICK
3	LABS PHS DIR IBTS	BARRY DOYLE	BARRY DOYLE
3	SSCD HEAD OF DEPT IBTS	AILEEN FARRELLY	AILEEN FARRELLY
4	QUALITY ASSURANCE REVIEWER IBTS	COLIN JOHNS	COLIN JOHNS
4	DIRECTOR OF QUALITY	KAREN BYRNE	KAREN BYRNE

Change Orders

Changes as described on Change Order: <u>Change Order No.</u>

Change Orders - Incorporated

Changes as described on Change Order: Change Order No.

IBTS/CO/0229/21

Valid on Day of Printing, Verify Current Version Printed By: Date:

IRISH BLOOD TRANSFUSION SERVICE PRODUCT MASTER FILE

TITLE: RED CELLS, SUITABLE FOR NEONATAL USE FOR

5 DAYS AFTER DATE DRAWN

Change Description:

Revise IBTS/PMF/SPEC/0203 to IBTS/PMF/SPEC/0212 and IBTS/PMF/SPEC/0232 to amend the product labels.

Reason for Change:

Fix to the labels with reference to IR 361/21/IBTS, IBTS/QA/PQ/0600 Deviation 012 and CC 134/21/IBTS

Change order No.:

IBTS/CO/0229/21

Referenced Documents

N/A

SmartSolve Roles

N/A

Training Type

N/A

SmartSolve Document Category

Category	Mobile	Cryobiology	Website	GDP
Yes / No	No	No	Yes	No

IRISH BLOOD TRANSFUSION SERVICE

PRODUCT MASTER FILE

Title: Red Cells, Suitable for Neonatal Use for 5 Days after Date Drawn

Name of Products: RED CELLS, Suitable for Neonatal Use for 5 Days after Date Drawn /

RED CELLS, Suitable for Neonatal Use for 5 Days after Date Drawn, Irradiated

E Progesa Codabar Component Codes: 54481 / 74421

E Progesa ISBT – 128 Component Codes: C7429V00 / C7442V00.

General Description: A red cell suspension obtained from whole blood by

centrifugation, removal of plasma with subsequent addition of a nutrient solution SAG-M. The removal of the majority of leucocytes is achieved by filtration. The selected donors meet

the additional criteria for neonatal use.

General Specification:

Parameter	Quality Requirements	Frequency of Control
Volume	231 – 355 ml	100%
Haematocrit	0.50 - 0.70 L/L	1%
Haemoglobin	≥ 40 g/unit	1%
Leucocyte Content	< 1 x 10 ⁶ /unit	1%
Haemolysis at end of shelf life	< 0.8% of red cell mass	4 per month (RED CELLS, 04333/E7429V00)
ABO Agglutinins	No HighTitre Anti-A or Anti-B	100%
CMV	CMV ab negative	100%

Labelling: See Appendix I

Storage: Red Cells, Suitable for Neonatal Use for 5 Days after Date Drawn

(including Irradiated) should be stored at $4^{\circ}\text{C} \pm 2^{\circ}\text{C}$.

Irradiation: Red Cells, Suitable for Neonatal Use for 5 Days after Date Drawn

may be irradiated up to 14 days from date of collection. Post

irradiation the storage is 14 days.

If irradiated, product code will change into codabar:74421and ISBT-

128 barcode: C7442V00.

Transportation: The air temperature of transport containers for units of

Red Cells, Suitable for Neonatal Use for 5 Days after Date Drawn (including Irradiated) should be maintained between 2°C and 10°C during transport from the Irish Blood Transfusion Service to the place

where they are intended for use. Transport time under these

conditions normally should not exceed 8 hours.

Indications for Use: Red Cells, Suitable for Neonatal Use for 5 Days after Date Drawn

(including Irradiated) are used for augmenting the oxygen carrying capacity of the blood where this is critically reduced in infants

requiring large volume transfusion, particularly in the surgical setting,

within 5 days from date drawn.

Precautions in Use:

• Compatibility of this component with the intended recipient must be verified by appropriate pre transfusion testing.

• Red Cells, Suitable for Neonatal Use for 5 Days after Date Drawn (including Irradiated) should be transfused through a

standard 170 – 200 µm filter.

• No solution should be added to the bag or to the giving set.

• Components should be inspected visually for defects, leakage,

abnormal colour or visible clots.

• Red Cells, Suitable for Neonatal Use for 5 Days after Date Drawn (including Irradiated) are not recommended in:

- large volume transfusion in neonates **unless** the red cells are

less than 5 days old.

- exchange transfusions in newborns.

Intrauterine Transfusions

This component does not contain platelets or soluble coagulation

factors.

Adverse Effects Include:

- Circulatory Overload.
- Haemolytic transfusion reaction;
- <u>Non-haemolytic transfusion reaction</u> (mainly chills, fever and urticaria)

• Pathogen transmission

- Despite careful donor selection and laboratory screening procedures, infections including Syphilis, Viral Hepatitis, HIV, HTLV 1 & 11 and other viruses and protozoa (e.g. malaria) may, in rare instances, occur.
- vCJD transmission
- Transmission of other pathogens that are not tested for or recognised.
- The risk of CMV transmission is minimal as the components are leucodepleted
- Sepsis due to bacterial contamination (reduced but not eliminated by bacterial screening)

• <u>Immunological effects</u>

- Alloimunisation to HLA, HPA and red cell antigens
- Post Transfusion purpura (PTP), especially in parous female recipients
- Graft versus host disease due to transfusion of viable lymphocytes can occur, but is minimised by exposure of the suspension to ionising radiation before transfusion
- Transfusion related Acute Lung injury (TRALI) by donor HLA/granulocyte antibodies

• Metabolic upset

- Citrate toxicity, especially in neonates and in patients with impaired hepatic function.
- \(\Tau^{+}\) in massive transfusions, especially where patient is hypothermic or acidotic or has impaired renal function.
- Hypocalcaemia.
- Hypoglycaemia.
- Hypokalaemia.

Iron overload

- In patients on chronic red cell transfusion support programmes.

Serious Adverse Reaction

Please inform the IBTS immediately about any event relating to suspected bacterial sepsis/ transfusion associated bacterial sepsis

Serious adverse reactions should be reported to:

National Haemovigilance Office

Irish Blood Transfusion Service National Blood Centre Quality Assurance Manager
Irish Blood Transfusion Service

ITHER
National Blood

AND

AT EITHER

James's Street Dublin 8

OR

Munster Regional Transfusion Centre St Finbarr's Hospital glas glas status Douglas Road, Cork

APPENDIX 1

E Progesa Codabar Component Code: 54481

E Progesa ISBT – 128 Component Code: C7429V00

Product NameRED CELLS, Suitable for Neonatal Use for 5 Days after Date Drawn

Shelf life 35 days

Labelling and Barcode:

(for illustration purposes only – barcodes not suitable for scanning – label not to scale)



IBTS ver 4.0

RED CELLS, Suitable for Neonatal Use for 5 days after Date Drawn

Store at 4°C ± 2°C



Drawn 05 May 2021



CMV Antibody Negative

Expiry 09 June 2021 23:59

C-c+E-e+K-HbS-Neg



This component must not be used if there are visible signs of deterioration. This component may transmit infection Must be administered using a suitable

Must be administered using a suitable transfusion set incorporating a 170 – 200 µm filter. Collected into CPD anticoagulant and suspended in 105ml of additive solution containing, in mmod/L NaCl 150, Clucose 45, Administ 125, Mannitol 29.



Expiry 09/06/2021



O Noa

APPENDIX II

E Progesa Codabar Component Code: 74421

E Progesa ISBT – 128 Component Code: C7442V00

Product Name RED CELLS, Suitable for Neonatal Use for 5 Days after Date Drawn, Irradiated Shelf life 14 days

Labelling and Barcode:

(for illustration purposes only – barcodes not suitable for scanning – label not to scale)



IBTS ver

RED CELLS, Suitable for Neonatal Use for 5 days after Date Drawn, Irradiated

Store at 4°C ± 2°C



Drawn 05 May 2021



IRRADIATED CMV Antibody Negative



This component must not be used if there are visible signs of deterioration. This component may transmit infection

component may transmit inrection
Must be administered using a suitable
transfusion set incorporating a
170 – 200 µm filter. Collected into
CPD anticogulant and suspended in 105ml
of additive solution containing, in mmol/l: NaCl 150, Glucose 45, Adenine 1.25, Mannitol 29.



Expiry 09 June 2021 23:59



C-c+E-e+K-HbS-Neg



74421



Expiry 09/06/2021

